

Applying Imidacloprid 2F (21.4% or 22.6% Flowable) by Foliar Spray

In general, SGH does not recommend applying hemlock treatment by foliar spray because:

- the concentration is very dilute and the level of protection is low,
- the residual protection period is very short and treatment must be repeated frequently,
- the cost on a per-inch-per-year basis is high, and
- much of the spray may miss its target and drift onto non-target plants or beneficial insects.

However, there are two cases in which foliar spray may be advisable:

- spraying a short hedge or a “nursery” of saplings where you can reach the whole plants, or
- spraying the lower branches of a tree that is also being treated by soil application to give those limbs relief sooner than later.

The best seasons for foliar spray are spring through fall. **Use only on a calm day in early morning or late evening** (the stillest times of the day) to avoid wind drift onto beneficial insects or the plants they visit and to avoid sun-scorch of the needles.

Step 1. To make a master batch of treatment mixture, follow the proportions of water and amount of product shown in the Mixing chart below. Put the water in the sprayer tank or mixing container first. Then add the corresponding amount of product and mix thoroughly.

(Hint: 1 fl. oz. = 2 tablespoons. 1 tablespoon = 3 teaspoons)

MIXING THE MASTER BATCH	
The label indicates using 1.5 fluid ounce (3 tablespoons) of product with 100 gallons of water. For smaller tanks, mix as follows:	
Water	Imidacloprid
50 gal.	1 tablespoon + 1 ½ teaspoons
25 gal	2 ¼ teaspoons
10 gal	1 teaspoon
5 gal	½ teaspoon

For a sprayer that holds less than 5 gallons, use 1/2 teaspoon of product with whatever amount of water the tank holds.

Step 2. Spray thoroughly on all parts of tree – stems and undersides of branches as well as top sides. Wear long sleeved shirt, long pants, shoes and socks, goggles and a mask while spraying.

Step 3. When you’re finished treating each tree, mark it to indicate it’s been treated. Give the treatment time to dry before children or pets are allowed in the area.

Step 4. Keep an eye on your trees for signs of reinfestation and repeat treatment as soon as the first new egg sacs are spotted.

CALL THE HEMLOCK HELP LINE 706-429-8010 WITH ANY QUESTIONS.

