

# Applying Imidacloprid 75 WSP (2.25 oz) by Foliar Spray

In general, SGH does not recommend applying hemlock treatment by foliar spray because:

- the concentration is very dilute and the level of protection is low,
- the residual protection period is very short and treatment must be repeated frequently,
- the cost on a per-inch-per-year basis is high, and
- much of the spray may miss its target and drift onto non-target plants or beneficial insects.

However, there are two cases in which foliar spray may be advisable:

- spraying a short hedge or a “nursery” of saplings where you can reach the whole plants, or
- spraying the lower branches of a tree that is being treated by soil application to give those limbs relief sooner than later.

The best seasons for foliar spray are spring and fall. Use in early morning or evening to avoid sun-scorch of the needles or harm to beneficial insects during the day. Also choose a calm day to avoid wind drift.

**Step 1.** To make a batch of solution, follow the proportions of water and packets of product shown in the Mixing chart below. Put the water in the sprayer tank or mixing container first. Then add the corresponding number of packets and mix thoroughly. Warm water works best for dissolving the packets.

<b>MIXING</b>		
Use one 2.25-ounce packet of product with 420 gallons of water. This yields a concentration of about .11 gram of active ingredient per gallon of solution. For smaller tanks, mix as follows:		
<b>Water</b>	<b>Imidacloprid by ounces</b>	<b>Imidacloprid by level measure</b>
300 gal	1.6 ounces	3 tablespoons + 1 teaspoon
200 gal.	1 ounce	2 tablespoons
100 gal.	0.5 ounce	1 tablespoons
50 gal.	0.25 ounce	1 1/2 teaspoons
25 gal.	0.125 ounce	3/4 teaspoons
10 gal.	0.05 ounce	1/3 teaspoon
5 gal.	0.025 ounce	1/6 teaspoon

For a sprayer that holds less than 5 gallons, use 1/6 teaspoon of product with whatever amount of water the tank holds.

**Step 2.** Spray thoroughly on all parts of tree – stems and undersides of branches as well as top sides. Wear long sleeved shirt, long pants, shoes and socks, goggles and a mask while spraying.

**Step 3.** When you’re finished treating each tree, mark it to indicate it’s been treated. Give the treatment time to dry before children or pets are allowed in the area.

**Step 4.** Keep an eye on your trees for signs of reinfestation and repeat treatment as soon as the first egg sac is spotted.

**CALL THE HEMLOCK HELP LINE 706-429-8010 WITH ANY QUESTIONS.**

